05030 Z

Scientific and Technical Information Center

SEARCH REQUEST FORM

Access DB# 83264 11A16

Requester's Full Name: Lynda Guo Examiner #: 79756 Date: 1230/02 Art Unit: 1657 Phone Number 30 605-1200 Serial Number: 10/070,018 Mail Box and Bldg Room Location 11801-mail Results Format Preferred (circle): PAPER DISK E-MAIL
TIAID - DIFFEE
If more than one search is submitted, please prioritize searches in order of need.
Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc. if sknown. Please attach a copy of the cover sheet, pertinent claims, and abstract.
Title of Invention: Novel target for antiparasitic agents and inhibitors thereof
Inventors (please provide full names):
Earliest Priority Filing Date: *For Sequence Searches Only* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the

Point of Contact: Mona Smith Technical Information Specialist CM1 6A01 Tel: 308-3278

STAFF USE ONLY	Type of Search	Vendors and cost where applicable
Searcher M. Boy 17H	NA Sequence (#)	STN
Searcher Phone #:	AA Sequence (#)	Dialog
Searcher Location.	Structure (#)	Questel Orbit
Date Searcher Picked Up. 1/2/03	Bibliographic	Dr.1.ink
Date Completed 1/17/03	Litigation	Lexis Nexis
Searcher Prep & Review Time & C.	Fulltext	Sequence Systems
Clencal Prep Time:	Patent Family	WWW/Internet
Online Time	Other	Other (specify)

```
81..W 1...E
  Bile IndiMEDINE Booleen-Look Jan WI
Bule of thisis Freviews Booleen-Lile Jan WL
                   Total Communication of the Com
                   44:8010earon B. Ditea Red Boo Devi-Off Cah Wi
  5.114
                              o sella imat ter gar info
                   n:Dissertation Aps Unline 1881-1883 1. 8
  2 114
                       :CAB Abstracts 1370-2170 Dec
20113 CAB International
                     l:ELSEWIER BIJBASE 1994-1718 Jan W2
                            N. 113 Elsevier Science B.V.
                   Tr:EMBASE 1874-1883 Jan Wi
Tr:EMBASE 1874-1883 Jan Wi
Tr:U175 Elsevier Science B.T.
M4:01887-EB148 1981-188 Mic Wi
  111
                  2. Us Capan Schence and Tech Corp 127
Mr. Seneral Sch Aps Full-Text 1954-1812 183
 File 144: T37 Health AWellness DB SW 147K-20. * Dec Wi
 5. 0003 The Gale Group

6.10 107:TowFile 1948-2000 Nov W3

7.10 107:TowFile 1948-2000 Nov W3

7.10 107:TowFile 1948-2003/Deb
                           (5) X003 CAB International
 File 1:5: Doslogical Record Online(R) 1978-2002/Dec
                           to 2002 BIOSIS
 File 340:CLAIMS(R)/US Patent 1950-03/Jan 14
                          (a) 2003 IFI/CLAIMS(R)
 File 381:Derwent WFI 1963-2002/UD, UM &UP=200303
                          3 2003 Thomson Derwent
 File 307:Derwent Biotech Res. 1982-2013/Jan W1
                          o .113 Thomson Forward & 181
Pule of H: Durrent Histeon Abs 1983-2002 Dec
 File 434:SpiSearch(R) Cited Ref Spi 1974-198 /*/wo
                          (a) 1998 Inst for Sai Info
 File 440: Current Contents Search R, 1990-2003 Jan 17
                        (a. 2003 Inst for Sci Info
 ? as
                                       Description (PARASIT? OR FUNG? OR BIOCID? OR INSECT? OR BACTER?) AND
365
                      Items
                     48955
                                SCREEN? OR ASSAY? OR TEST?: AND "PHOSPHATE? OR PHOSPHATASE?

1 AND ENCYME? S. INHIB? OR SUPPRESS?:

SI AND ENCYME? S. "ACTIVE OR INHIB? U.B. SUPPRESS?.

S3 AND UWORM? OR PROTOSOA? OF NEMATUDE? OR MITE?
                   1:477.
1/44.
. . .
                           i -- i
                                       FD unique items
                           TO A FULL ARTIQUE TOERS

11 SE AND STREHALOSE W.6 P. SLYSEPTE W.3 OF MAINVITCH W.1 P.2-

FRITCH W.6, W. PHOSPHATER,
- Fl - unique items
                                 liter i frim file: f
IIAN FR File E:Firsis Fremiews F
 1. -177. 4 - Pindia W ... 21,111024778
April to analyging or inharitary of Trypania na price, prognizionerate
```

- Kinaret Kiştidat'ı nılı a nüvel bindingir biliti a nemin emleşirette yazı Caten Sine.

AVIH.B: Press: Jerire J; Thie Jungwil; Holyn Medinia I; Furnher Freier. R ; Van Vidinis Wesley J; Verlinde Jhristigne I M J; Hol Wir J J a ; holy M. thae. H a

WVIH & ANIBERVE of Department of Themserry, University of Washington, Charleston, WA, 2018; for the new washington, automata

o defiale du Lenge de Medermal Chemieney de la 1941-1941 (Mintenges L.,

MROOME PEER 1990: LL-LEUP 1990: MROO OFFE: Artible BEO.FO OFFE: Abstract LADSTAFF: English NOOMARY LADSTAGE: English

Abulbant: As part of a project aimed at structure-based design of adenualne abaliques as drugs against African trypanosomiasis, No-, D-amino-NV-, and In -substitution agents including we have synthesize i and the specific estarion structure- activity relationships for inhibiting Trypan, soma it well plysteimal phosphoglyperate kinase FGM, glyperaldenyde-3-phosphate denydrogenase GAPDH, and glyperol - 3 - phosphate whydrogenase GPDH: Evaluation of M-ray structures of parasite FGM, GAFFH, and GFDH complexed with their adenceyl-bearing substrates led us to generate a series of adenosine analogues which would target all three Fnnymes simultaneously. There was a modest preference by PGK for Me-substituted analogues bearing the 2-amino group. The best compound in this series, l-amino-Né-,l"- p-hydroxyphenyl, ethyl -adenosine (40), ausplayed a 23-fold improvement over adenosine with an 1080 of 131 hum. $2-\sqrt{2}^{-1}$ p-Hydroxyphenyl, ethyl) amino, adenosine (465) was a weak inhibitor of T. brucei PGK with an ICSO of 500 muM. To explore the potential of an additive effect that having the N6 and N2 substitutions in one molecule might provide, the best ligands from the two series were incorporated into N6, N2-disubstituted adenosine analogues to yield NG- 1"-phenylethyl: -2-;(2"-phenylethyl;aming-adenosine (63) as a 31 max unalret roof T. prusel PSK which is 170-full more potent than the submissible template. In contrast, these series gave no compounds that unmikited parasitic GAFDH or GPDH more than 10-20 when tested at 1.1 mM. A 3.1 ANG M-ray structure of a T. brucei PGM/48b complex revealed a binding mode in which the nucleoside analogue was flipped and the ribosyl moiety adopted a syn conformation as compared with the previously determined binding mode of ADP. Molecular docking emperiments using AMP and SAS program suites reproduced this "flitted and rotated" binding ∷ade.

AB A ltem 2 from file: 8, TIAL:3 P File | 5:Biosis Previews(P | 1 113 HIOSIS. All rts. reserv.

PRIVATE Austrant

"THAT BIDGE NO.: TILLESTAGES
AN IMPRITED PURIFICATION OF SEVENOMES FROM THE EPOTYPHIC TRYSOMANTIS TEN
BUTTER AMAI BOA; WANG OUT
AUTH BOALLEST, MEL. MICROBINE, CTANFORD UNIO, JUNE, MEL., JUANFOR,
TA.
TOPNAL: MIL BIOCHEM PARASITOL 21 (3), 1984, JULI-WILL 1984
FULL TOPNAL NAME: Molecular and Biothemical Barawittlery
TIEM: MRIPL

LANCER ENGLISH

About A tip The gly o stoke of in vita, or which by the tripper activity is now It live at some prices were purified by three different procedures and the restable to practical by selection problems by plant security of essays can be as in or ar types platemy lyamylamide gelelettriphlievis. Ventringstripti : Verrei imuni Perivili gradient filliwei by Asim se draifent Mintriculation resulter in the least enricher guyo's has preparation. Ministratation on a pre-comed Myodueno oraquent gave an improved preparation for the bost nomogeneous preparation in interpolityons nes was it lutera atter mentritugation on two sundessive sum se gradients. Regressions publified by buth the Mybrakho and souble submise gradient principles appeared larger than in situ glyphsomes presimably are to an om the effect resulting from disruption of the granular matrix of the organelles. Mevertheless, there appears to be no loss or disternal contents due to the swelling of the organelles. The objections of the fulla stream form trypomastigates purified by the same protection of or r wever, it sign is Swelling. A comparison of slyng new jurished time promy violatry pomastigates and plucistream from wrypomastigates prepared ly the same diuble subrise pribedure demonstrated that in the physisime of proposition trypomastiques: I pactivities of new kinase, phisphogluorse isomerase, phosphofruotose kinase, aliolase and in sincipy erate kinase and diminished by -1-1/10, up a missings of ilymeraldehyde-d- phosphate dehydrogenase, triose phosphate is merase and ilymeral - 3 - phosphate dehydrogenase remain un manged r are unly slightly reduced; 3, there is an appearance of four major new proteins, aming which could be phosphoenol pyruvate carpiwykinase and malate dehydrogenase. These observations are in basic agreement with those by Hart et al. Mol. Biochem. Parasitol . 12, 25-38, 1984;.

1356

LAN WA HE: ENGLISH

TVABV3 | Titem 3 from file: 5; DTALUG.RUFile | 5:Biosis Previews(R, 5 .003 BIOSIS. All rts. reserv.

ISALIBBIO BIOSIS NO.: 000074019468

1. TALIDATION OF MALATE DEHYDROGENASE EC-1.1.1.30 ALEMYLATE KINASE
EU-2.7.4.3 AND GLYCOLYTIC ENZYMES IN GLYCOSOMES AND THE THREONINE PATHWAY
IN THE MITOCHONDRION OF CULTURED PROCYCLIC TRYPOMASTIGCTES OF
TRYPANOSOMA-BRUCEI
AUTHOR: OFFERDOES F R; MARKOS A; STEIGER F F
AUTHOR ADDRESS: RESEARCH UNIT FOR TROPICAL DISEASES, INTERNATIONAL INST. F
CELLULAR AND MOLECULAR PATHOLOGY, IOP, AVENUE HIPPOGRATE 74, B-1/01
BRUSSELS, BRIGIUM.
UNFNAU: MOL BIOCHEM PARASITOL 4 (B-6). 1941 [RECD. 1967 . 291-910. 1961
FULL J URNAL MARK: Molecular and Biochemical Farasitology
CEN: MBIPO
FEO RO TYPE: Abstract

ARSTRACT: Prohydlic dulture forms of the human and dattle parasite T. nradel stock 427 were screened for the presence of enhymes involved in glybolysis, mitochondrial energy metabolism and threshine degradation. The enhyme activities in the prohydlics were obspaced with those of the closesteam forms The specific activities of glybolytic enhymes represented for the respective levels in the plocastream form, except for nextkinase [EC 2.7.1.1] which was limited required. Fellower that how we that the enhymes involves in the garly dequate of the slybolytic pathway, i.e., from nextkinase to propped any relate single Fig. 3.1.1.1 and the enhymes NAL--linear slybolite or expectate.

denyar genase [RO 1.1.1.0] and glyceril minase were all present in glyceshes equilibrating at a density of 1.00 y only in substance granulars. Malate denyar genase was firstly more active in procyclics than in collectron firms. This increase in approximative was the result of the appearance of malate denydrogenase in the alyceshes of the procyclics, in addition to mitionomyrial and beliesappearties which were present the first stades of the life cycle. Slyceshes gentalized part of the applicable dinase activity, which was also associated with the minimum of the denydrogenase [RO 1.1.8s.), a detnor with the mitionomyrial denydrogenase [RO 1.1.8s.), were located in the mitionomyrial which had a density in sucrese ranging from 1.10-1.1-3 cms. This organelle also contained 1-threchine F-denydrogenase [RO 1.1.1.1.1.0] and carnitine acetyloransiciase [RO 1.1.1.1.0] and carnitine acetyloransiciase [RO 1.1.1.0] and carnitine [RO 1.1.0] an

1941

T AB 4 (Item 4 from file: 5)
LIALOS R(File | 5:Biosis Previews(R)
| 3 | 2003 BIOSIS. All rts. reserv.

ICHITHGE BIOSIS NO.: 000069026084

TREHALOSE 6 PHOSPHATE SYNTHASE EC-2.4.1.15 FROM
DICTYOSTELIUM-DISCOLDEUM PARTIAL PURIFICATION AND CHARACTERIZATION OF THE ENDYME FROM YOUNG SCROCARPS
AUTHOR: KILLICK K A
AUTHOR ADDRESS: DEP. DEV. BIOL., BOSTON BIOMED. RES. INST., BOSTON, MASS. 02114, USA.
UCUFNAL: ARCH BIOCHEM BIOPHYS 196 (11. 1979, 121-133, 1979)
FULL COURNAL NAME: Archives of Biochemistry and Biophysics
CUDEN: ABBIA
RECURD TYPE: Abstract
LANGUAGE: ENGLISE

ABSTRACT: Trehalose 6 - phosphate synthase was solubilized from young sproparps of the cellular slime mold, D. discoideum, by a freeze-thaw cycle and was subsequently purified about 160-fold using streptomycin sulfate precipitation, (NH4)2SO4 fractionation, DEAE-cellulose chromatography, heat treatment in the presence of heparin and molecular sieve chromatography on columns of Bio-Gel A-1.5 m. The purified enzyme was maximally active at pH 6.5, showed an absolute specificity for G-6-P as glucosyl acceptor and a relative specificity for the glucosyl donor in the order: UDF-glucose, GDF-glucose and ADF-glucose. Although hoparin and chondroitin sulfate activated the synthase, the order of gludsyl donor specificity was not affected. Sther activators of trehalose 6 - phosphate synthase were KCL, Mg2+ and EDTA, while detergents had little effect. Although synthase activity was reduced f to BIT upon the omission of Mg2+ from the Lassay mixture, an absolute dependency for Mg2+ could not be demonstrated. Evaluation of the apparent Embralues for partially purified synthase preparations demonstrated that for each of the synthase substrates, the lineweaver-Burk plots displayed ourplex cimodal kinetics. Estimation of the Em after extrapolation of the viral mit line purtions of these plots yielded values of 1.2 and 3.1 mm Web-F and 1.5 and 3.2 mm VDF-glubose. Comparison of the latter parameters with the religiar levels of TDF-glusise and G-6-F in Distyratelism supplies that it the orserved bimodal kinetics are the consequence of ruligic kinetically distinct forms of the symmask, the activation of

```
tional to symplectic during since \tau is further. In ( ) is provide a fact have for the process of these is symmet.
```

Abstract: A number of chemotherapeutic compounds effective against 1. prunel were tested as inhibitors of the purified glycerophosphate wides and suramin were potent inhibitors of the dehydrogenase components this multilenayme complex. Suramin is a potent competitive inhibitor to the twidese with a Mi of 4.1 .mu.M with respect to glycerophosphate. The Mm for glycerophosphate for the encyme decreased from 6.5 to 1.7 mm in the presence of bovine serum albumin while the Vmax was increased 2-to 3-fold. Human and bovine serum albumin can protect the oxidase from inhibition by suramin, by preferential binding of the drug. Analogs of suramin with little or no chemotherapeutic value are less effective inhibitors of the oxidase, and the correlation between therapeutic action and potency as inhibitors sudgests that this encyme is 1 in the principal sites of action of suramin in vivo.

19**

That is a litem of from file: 5.

.

ITALUG R.File | 5:Bicsis Previews.R)
5 1003 BIOSIS. All rts. reserv.

"219-ER1 BIOSIS NO.: 000064041070
A BACTUMETRIO ASSAY FOR TREHALOSE & FH.OPHATE SYNTHETASE F0-2.4.0.15
AUTHOR: MILLION K A
100FNAL: ANAL BIOCHEM 79 (1-2). 1977 310-314. 1977
FULL CUTENAL NAME: Analytical Biochemistry
0.0EN: ANFOA
FFO.EL TYPE: Abstract

spring intry, of other encyme pesture the synthetias cratalynes to synthesis of firehalfseries, greater sensitivity at 1 w levels of product; and invents livity to competing size reactions and invents of interfere with the firehalf of assay of each, polyric high time time prospering assay synthesis of polyrication of nucleous appropriates.

- Who is them to rive rive; at CIAN to be blue of the terms. Win Also bull-Text To one to the HW Wilson Mi. All (ris. reserv.

ABSTRANT: The inree-dimensional structures in tryptopnan synthase, far.amoly. Phosphate synthetase, glutamine phosphoriposylpyrophosphate amiastransferase, and asparagine synthetase have revealed the relative lightions of multiple active sites within these proteins. In all of these polyfunctional encymes, a product formed from the datalytic reaction at the active site is a substrate for an encymatic reaction at a distaluative site. Reaction intermediates are translocated from one active site to the next through the participation of an intermolecular tunnel. The function in tryptophan synthase is (similar)2b. A in length, whereas the tunnel in tryptophan synthase is (similar)2b. A in length, whereas the tunnel in carbamoyl phosphate synthetase is nearly 101. A fond. Kinetic studies have demonstrated that the individual reactions are coordinated that up allosteric coupling of one active site with another. The factors active active site with another. The factors active from coming in contact with the external medium. Reprinted by persission of the publisher.

[4118282 H.W. WILSON RECORD NUMBER: BGS199118282 Emperimental evolution and its role in evolutionary physiology. Hemsets, Albert F. Lenski, Richard E. American Coologist (Am. Cool) v. 39 no2 [Apr. 1984, §. 846-62] URBITAL FEATURES: bikl il | 1880; 1003-1169 LANGUARE: Frylish | 1007FF (F. FUBLICATION: United States W. B. 1007F. 1768)

ABCTBACT: Four general approaches to the story of evolutionary physics by epitementically-based companies as, sometic analyses and manipulations, phenotypic plasticity and manipulation, and selection of the example of the latter, the application of laboratory selection experiments to the story of a selection experiments to the story of a selection selection, aliteratives in adaptive pattern of the full stars and specialists. A close of the harterion Example of the full cases of was replicated.

The trip plantage and allowed to reproduce to 1., the personnels in a variable theorem and convictment alternating retwien a land 4. Heaventh As five the 1. The 1.7, intress and efficiency of results and extrapolations, as an attress twenternal and extra any organism of the direction of the first and efficiency in the other intrapolation and extra any organism of the direction of the first ase, has an attress and the province a variable efficiency in the second of the first and the first

Abor Item 1 irom file: 149 CIAL NE File 149:T37 HealthWellness IB om The Sale Group. All rts. reserv.

ultions SUFFILES NUMBER: 94071081 Under SMAT Tils 4 E E Butt Temp Thanges in membrane polar lipid fatty spiks of sessione paspalum in thepomeento liw temperature emposure. Turninass Oriente . Typol, J., Euwell, S.L., Lundan, B.E., Baird, W.U. Trop Somether, 40, 6, 2031 To

FUBLICATION ELEMAT: Magazine Journal; Bereseva (1820) (11-14-W) LANGUAGE: English (RECORD TYPE: Fulltent; Abstract (1ABORT AUTIENTE) Abademio; Trade WORD COUNT: (8317) LINE COUNT: (1881)

ANTHUR ABSTRACT: Seashore paspalum (Paspalum vaginatum Sw. is a warm-season turigrass, best known for its superior salt tolerance. Plants are subject to injury during winter conditions along the northern boundary of their cone of adaptation. New cultivars that are more telerant to lew temperatures are needed for use in the transition zone. Cold tolerance has been correlated with the degree of unsaturation in membrane lipid fatty asids. The attracted factly asids are thought to aid in maintaining membranes in a fluid state necessary for biological functioning 'nemecohasic a imposition of membrane lipids in three genotypes differing in cold tolerance. A second objective was to investigate changes in fatty acid content in these genotypes during exposure to low temperatures. Cold-treated plants were emposed to a 10-n photoperiod at t degrees)/14(degrees)C day/hight temperatures and light intensity of 25% miori mil m.sup.-1 s.sup.-1) photosynthetic photon flum density for 3 wk. Bhicomes and prowns were harvested at 7-d intervals. Total livids were extracted and the polar lipids separated by thin-layer chromatography. Fatty acids were identified by gas chromatography [32] and mass spectroscopy. In all three genotypes, the two saturated ratty arids, palmitid adid and stearid adid, did not thange during told treatment. Introungaturated singlened add increased signorisantly during low temperature exposure. The magnitude of change was greater in the concernees and more cold-tolerant Fi Sixila-I - Gealsiel', than in the concerned stely cold-tolerant. Adalayd' or in the cold-susceptible, parse-textured FI 198942. These findings suggest that accompliation or linglenia atld partly explains the differential response on their abid . Teranie i

The AP 17 Item I from tile: 381 ITAL SP File 381:Terwent WEI The File The rest Franchi, All rts. sees ro.

```
Assessini imminite linninitirs of sugar sider aloon loopiisphatases
         I do not not do includes candidantificada o toppolos beasuring the compute
 If the administrates and antifultation of process of and antifulty of the process of the daministration of the contract of the daministration of the contract 
                                    Find Tage Application All 17 17 Find Bill BR. 128 1
                                                                                                                                       Min: late
                                                                                                                                      A literater Links &
  Including Applications of Type Cate of HP endie 5 Adjenning
  Estent Details:
Estent Du Fins Lam Pa Main 180 - Filims Notes
Est 18-18: Al Folk Millell 41
 Asstrant Basin:
              N VELTY - Assessing substances as direct inning to inning the prospect of side sugar alounce phosphatases on golds where the sugar subar at the sugar substances of abnormal conditions comprises:
                          a. Measuring the enzyme activity in the presence of the
               inhabitar and sugar-sugar alcahol phasphate in a biological medium;
                          b Determining that the substance is an inhibitor if the encyme
                 activity is reduced.

DETAILED DESCRIPTION - An INDEPENDENT CLAIM is also included for an
               insibitor obtained by the method of the invention.
                          USE - The method is used to isolate inhibitors or sugar/sugar
              alruhil phosphatases and lead to a cynotoxic build up of an
              invermediate normally produced under conditions of stress claimed .
              The inhibitor can be used as a bicoide along with an anticundal abent
              e.:. an abble blaided .
                      ALVANTAGE - The inhibitors have a novel be manish of a mich
              compared to known anti- fungals and would therefore be less likely to
             encounter resistance among pathogenia fundi . Known antifundals of on
             have undesirable side-effects e.g. towidity to mammalian calls which
             would not be encountered with the new innihitors.
                         pp; 28 DwgNc 0/9
   7 %AB 11
                                            Item 2 from file: 351,
 DIALDS (R) File (351: Derwent WPI
    d 2003 Thomson Derwent. All rts. resert.
 WBI WAY NO: 0000-288229/200124
 Relateb WPI Abn No: 2011-246803
WBAM Ann No: 001-100481
     New innihiters and spreening assay for innihiters or suppressins if
Transport of the spharages of rangar prospectases. Farent Assigned the TETTEN SEC. & LETTEN SEC. & L
 Batent Family:
                                          Harver W
W . Miller
Am . Miller
                                                                                                                                         Final Late
                                                                                                                                                                                        West K
```

```
Enclosity Applications of the Type Care in MECO one of the Action, with Albertain
Parkir Ikrailk:
Estant Domition Fig. Est. Main 1870 - Fillins 1986
World Tierro A. Est a Ti., Fill 4.
     Figure 1 of the Control of the Contr
      il 17 di 18 Marin Francisk (T
Austrach Basin: W. L. 116gb Al
Arstran Basic:
               | November - A streeming assay it inhthit re it suppressies
          t sugar alrehel phosphatases or sugar phosphatases using encymes
        t dus troitings, kapteria, insects, hematiques, mates de
       products , is now.

IFTAILED DESCRIPTION - A mother for assessing the activity of test substances as inhibitors of a first bell encyme converting a sugar phosphate into a sugar, or a sugar alcohol phosphate into a sugar or sign alcohol is accumulated in sugar alcohol in bells. The sugar or sign alcohol is accumulated in
        large quantities by the cells, e.g. under conditions deviating in mother timel growth condition of the target cells or as a reaction to stress
        conditions, the inhibition being directly of the first encyme or
        indirectly. The method comprises:
               a contacting a test compound with a biological medium
        simprising the sugar phosphate or sugar albenol phosphate and the
        first onsyme;
                or, measuring the lactivity in the medium which depends on the
        e^{i\pi^2\pi^2} by of the first oneyme ;
                or repeating steps (1, and (2, with turther test compounds; and (d) selecting at least 1 compound which reduces activity of the
       ensyme compared with the same medium without the inhibitor; and elegationally assessing the activity of a second cell ensyme
       which is involved in the synthesis of the corresponding sugar
       phisphate or sugar alcohol phosphate .
               The selecting step includes selection of linhibitors which region
       the activity of the first engyme while maintaining a viable
       a minity of the sessing endymes.
                INDEFENDENT CLAIMS are included for the collowing:
                      inhibitors identified by the above meaning
                  x preparations comprising 1 or more of the inhibitors;
                      f a biscide acting on fundit, threets, hematides,
       capterial or other organisms assumulating large quantities of sager
       a, the large sugar in response to stress, comprising the inhibiture,
      4 a method of increasing the sugar phosphate or sugar along phosphate consumar along phosphate content in a target cell, particularly a manualian parameter, comprising using an inhibitor to require the according of a tirst
       rell lengymé converting a sugar (phosphate into a gugar or sugar
aconologicosphate into a sugar alcoholy and
                       alrethol of reducing of impairing the paramphicity of a
      numbellar parasite by promoting myper-arrumulation to a sugar
                             or a wigar almost (programe).
              A TIVITY - Antiparasiting of industries ; cartering a ; protocolor
           DE THAUTOM TE ADMITTU - Translose - é - poioposte o primposante.
```

The inning this.

Various to bentrate his of test of mpounds were applied to
Various types terevisiae, and from their inning try activity the fit
was factuated. Test ormgrund II had an I'm or solarly to the power

The state of the second of the proventing parasition, particularly funds, , raterial or
proventing parasition, particularly funds, , raterial or
proventing the time, or a mematide , line to , with or more
intestable, in a number, animal or plant.
pp. 1 # Twill out.

Guo 10 270, 31: Page 1

Copyright of the articles to which research is database refer to held by the publishers listed in the Foundation of PB field available for records published or updated in Character Wistracts after Desember 26, 1996, unless otherwise indicated on the adjunct publications. The CA Lexicon is the copyrighted interesting property of the the American Chemical Scotlety and is provided in 12 assist you in searching databases on STN. Any dissemination, a few controls, copying, or storing of this information, without the prior will endonsent of CAS, is strictly prohibited.

FILE COVERS 1907 - 17 Jan 2003 VOL 13- 10. 4 FILE LAST UPDATED: 16 Jan 2003 (2003) 11- 18

This file contains CAS Registry Number of a say and accurate substance identification.

```
=> d stat que
                            39 SEA FILE=FEGISTRY TREHAL . F-+-EHOSPHATE?/CN
                           39 SEA FILE=FEGISTRY INERAL CONTROL OF THOSPHALE? ON 195 SEA FILE=FEGISTRY GLYCH: 1-1-HOSPHATE?/CN 23 SEA FILE=FEGISTRY MANNICLE-HOSPHATE?/CN 5 SEA FILE=FEGISTRY SORECTION 1-4-PHOSPHATE?/CN 1 SEA FILE=FEGISTRY ERYTHSCOR
L3
14
L6
L7
                      18 SEA FILE=FEGISTRY SUGAN AND APHATERYON
325 SEA FILE=HOAPLUS L1 OR TARRAIL DE(W) 6 (W) PHOSPHATER
5865 SEA FILE=HOAPLUS L2 OR FLYTER L(W) 3 (W) PHOSPHATER
231 SEA FILE=HOAPLUS L3 OP TARRAIL DE(W) 1 (W) PHOSPHATER
112 SEA FILE=HOAPLUS L4 OR A FRANCIA DE W) PHOSPHATER
2 SEA FILE=HOAPLUS ARABIT DEC WARROUTH OR L6
3083 SEA FILE=HOAPLUS ERYTHFOLD WAR 4 (W) PHOSPHATER
69967 SEA FILE=HOAPLUS (?PARACITE F FUNG? OR BIOCID? OR ?INSECT?)
AND (SCREEN? OR ASSAURE DEC WORLD) FUNGE OR BIOCID? OR ?INSECT?)
18
L9
L11
112
L13
L14
L15
                      AND (SCREEN? OR ASSAY? FROM PHOSPHATASE? OR SUGAR(W) PHOSPHATAS
L16
                             E? OR L8 OR L9 OR L10 'F 11. "F L12 OR L13 OR L14
L18
                             52 SEA FILE=HCAPLUS L17 AND FOUNDER?
L19
                             15 SEA FILE=HCAPLUS L18 ALL L DM. OR BACTER? OR PROTOZOA? OR
                                   NEMATODE? OF MITE?.
```

=> d ibib abs mitrm 119 1-18

L19 ANSWER 1 CF 15 HCAPLUS COPYRIGH. ANS
ACCESSION NUMBER: 2002:695798 HOWER UP
DOCUMENT NUMBER: 137:226941

TITLE: Use of pertail to this for treatment of a number of conditions in the list for the l

```
SGURCE:
                                  POT Int. App...
                                   CODEN: PINWI.
 DOOTMENT TYPE:
                                  Patent
 LANGUAGE:
                                   English
 FAMILY ACC. NUM. COUNT:
     PATENT INFORMATION:
PRIGRITY APPLN. INFO.:
                                                              1-340054F F 20011101
1-335015F F 20011105
1-343523P P 20011220
OTHER SOURCE(S): MARPAT 137:227 - T
AB The invention relates to the use the part of the delayed effects of radiation therapy. Compds. that can be used the invention include methyl-2,3,4-trihydroxy-1-O-(7,17-3) & androst-5-ene-3.beta.-yl)-.beta.-D-glucopyranosid ronate. Formulation the steroids are also
       exemplified.
      9075-65-4, Glycerophosphate dehyd: : RESU (Biological study) (steroid hormone induction or the study) of mitochondrial GDPH and cytosolic malic enzyme in ration of the study; synthetic prepr. and use of certain steroids for treasure of a no. of conditions including
          blood cell deficiencies)
REFERENCE COUNT: 13 THERE AND CLIED REFERENCES AVAILABLE FOR THIS RECORD. A ATTOMS AVAILABLE IN THE RE FORMAT
L19 ANSWER 2 OF 15 HCAPLUS COPYRIGH. AUG
ACCESSION NUMBER: 2002:89878 Howard
DOCUMENT NUMBER:
                                136:156403
TITLE:
                                Methods for immorphism therapeutic targets for
                                treating inferrior disease
Shepard, Michael C.; Lackey, David B.; Cathers, Brian
INVENTOR(S):
E.; Sergeeva, Mile Comparison of the Extra ASSIGNEE (S): Newbiotics, Discount of the Extra Assignment (S):
                                 POT Int. Appl., 18. CODEN: PIMMIZ
SOURCE:
                                  Patent
LANGUAGE:
                                 English
FAMILY ACC. NUM. COUNT:
PATENT INFORMATION:
      PATENT NO. KIND DATE CONTINUE CATE
```

```
Gus 11 07., 018 Page 3
            W0 2002107780
PRICEITY APPIN. INFO.:
       This invention provides methods and thems to identify enzymes that act as enzyme-catalyzed therapy that act as enzyme-catalyzed therapy that activators and the enzymes identified by these methods. The provided by this invention are compds. Activated by the enzymes as well as
AΒ
       compns. contq. these compas.
       37250-69-4
       RL: BSU (Biological study, unclastic : ; 'AT (Catalyst use; THU (Therapeutic use); BIOL (Biological : ; USES (Uses) (identifying intrinsic enzyments), a therapeutic activators as targets for treat : ; is attooned dispase
            activators as targets for treating theathous disease
ŢΤ
       9025-72-3, E.C. 3.1.3.12
       RL: CAT (Catalyst use); PRP (Prop. ...; THU (Therapeutic use); BIOL
       (Biological study); USES (Tses)

(identifying intrinsic enzyme of therapeutic activators as targets for treated the sticus disease)
L19 ANSWER 3 OF 15 HCAPLUS COPYRIGH:
ACCESSION NUMBER: 2001:845255 AMA C
DOCUMENT NUMBER:
                                    136:34648
                                   Benes, enzymes, soled intermediates, and methods for all of mevalonate-independent isoprenoid by is pathway Adam, Petra; but all the pathway Fellermeier, Koron, Hebrit, Stefan; Fondich, Felix;
TITLE:
INVENTOR(S):
                                    Schuhr, Christ prom.; Wungsintaweekul, Juraithip;
                                    Jenk, Meinhart H.
                              Germany
PATENT ASSIGNEE(S):
                                    Ger. Offen., help.
SCURCE:
                                    CODEN: GWKKBK
DOCUMENT TYPE:
                                   Patent
LANGUAGE:
                                    German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:
```

```
LE, DK, ES, FI, FR, GB, TV, LOT, MO, NL, FT, CE, TF, BF, BC, OF, OG, OI, OM, GA, TV, LOT, MF, NE, SN, TO, TG PRIGRITE APPLN. INFO.: -10027821 A COIL ONG
   AB The present invention concerns enzymes . I intermediates of the
                     nevalonate-independent isopremont to select particles of the selection of inhibitor-resistant. In the selection of inhibitor-resistant.
                    consern genes coding for the enzymes . For innicitor-resistant variants of the enzymes, vectors which contain the vectors, and place to contain the vectors, and place to contain the vectors, and place to contain the medicine. Thus, the Bacillus subtilis and Escherichic to the medicinete-independent isoprencial biosynthes. The DMP synthase and DMP reductor to enzymes were used to prep. [U-1308]-20-methyl-D-erythritol-4
                     phosphate. The gene ygiE l-mermy- - se-t-phosphate synthase, gene yaeM l-mecmy-D-mylulose-E-ph synthase were used in preprinced and phosphosytidyl-2C-methyl-D-erm synthase were used in preprinced ygbP, i.e., gapE, lytB, ydeE, and so synthase were used in preprinced ygbP, i.e., gapE, lytB, ydeE, and so synthase were used in preprinced ygbP, i.e., gapE, lytB, ydeE, and so synthase were used in preprinced ygbP, i.e., gapE, lytB, ydeE, and so synthase were used in preprinced ygbP, i.e., gapE, lytB, ydeE, and so synthase in preprinced ygbP, i.e., gapE, lytB, ydeE, and so synthase or for preprinced ygbP, i.e., gapE, lytB, ydeE, and so synthase or for preprinced ygbP.
                      intermediates in the pathway.
  ACCESSION NUMBER: 2001:816926 H W COMBER: 2001:816926 
   L19 ANSWER 4 OF 15 HCAPLUS COPYRIGH:
                                                                                                 Structure of a real about dayl methylerythritol synthetase in a real and the rational design of
   TITLE:
                                                                                                   effectors
  INVENTOR(S):
                                                                                                 Noel, Joseph E.; a sman, Marianne E.; Richard,
                                                                                                 Stephane
                                                                                        The Salk Institute to r Biological Studies, USA
  PATENT ASSIGNEE(S):
                                                                                               POT Int. Appl., Prig. CODEN: PIMMO?
  SCURCE:
 DOCUMENT TYPE:
                                                                                                Patent
  LANGUAGE:
                                                                                                 English
  FAMILY ACC. NUM. COUNT:
  PATENT INFORMATION:
                    PATENT NO. KIND DATE - MICATION NO. DATE

      WC 2001033769
      A2 20011108

      WC 2001033769
      A3 20020829

                                                                                                                                                                       U 0001-US14371 20010503
The present invention provides the fixe of the enzyme

4-diphosphopytidyl-2-0-methylery the Synthase, a member of the syttidyltransferase family of enzymes. F-MF is a prot.
```

AB

```
intermediate in the mevalonate-in: \{0,0,1\} pathway for isogrenoid biosynthesis in a no. of proparty: \{0,1\} sms, in algae, in the plastids
                of plants, and in the malaria parasite. Cince vertebrates
                synthesize isoprenoid precursors and a newalchate pathway, our-me
              synthase and other enzymes of the structure-cased lesign of selective antipations.

Accordingly, the present invent:

Compass that inhibit enzymes of the structure-cased compass that inhibit enzymes of the pathway and pharmaceutical compass that inhibit enzymes of the pathway and bacterial terpencials:

Somethics for the structure-cased compass that inhibit enzymes of the pathway and bacterial terpencials:

Compassion as synthasis of the structure-cased compassion and methods for the pathway and bacterial terpencials:

Compassion as synthasis of the structure-cased compassion and methods for the structure-cased compassion compassion and compassion compassion
                treating a subject suffering from . bacterial infection.
 119 ANSWER 8 OF 18 HOAPLUS COPYRIGHT
 ACCESSION NUMBER: 2001:168177 % % . DOCUMENT NUMBER: 134:217175
 DOCUMENT NUMBER: TITLE:
                                                                            Sugar alcohol
                                                                           Sugar alcohol cases or sugar phosphatases a targets for
                                                                            antiparasitic and use of the inhibitors
                                                                            in blocides and consequipals
 INVENTOR(S):
                                                                     Thevelein, Ither or Dijok, Patrick K.U. Leuven Feld or & Development, Belg. PCT Int. Appl., 1985.
 PATENT ASSIGNEE (S):
 SOURCE:
                                                                            CODEN: PIXMDZ
 DOCUMENT TYPE:
                                                                            Patent
 LANGUAGE:
                                                                             English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:
               WO 2001016357 A3 20011129

W: AE, AG, AL, AM, AT, AU, A., A., EB, BG, BR, BY, BM, CA, CH, CN, CR, CU, CZ, DE, DR, DM, L., ., .3, FI, GB, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, CM, MM, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TU, CM, LR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, FW, C, CT, TU, TM

EW: GH, GM, KE, LS, MW, MZ, SI, CM, CM, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GB, C, CT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, C, CR, NE, NE, SN, TD, TG

EP 1081232 A1 20010307 B1999-202605 19990830

E: AT, BE, CH, DE, DH, ES, FF, C, CR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO

EP 108568 A2 20020522 CAR INC. SA 20000329
                                                                                                                                  114 11110-964054 20000929 Mg, PT, 127 Mg, PT,
                EP 1206568 A2 20020522
                           R: AT, BE, CH, DE, DK, ES, FR,
                                     IE, SI, LT, LV, FL, RO, NO.
                                                                                                                                       PRIORITY APPLN. INFO.:
              The use of an enzyme found in fung:, bacteria
               , insects, nematodes, worms, mites
               , protozoa etc. as a target in a screening
              assay is described by means of which is the capable of inhibiting
              the function of that enzyme may keep tied. The screening assay may include complete or purifiedenzyme assays. In particular, the properties.
```

```
relates to a screening assay for
                 suppressors of sugar alo, phospholoses sugar phosphatases, and more in particular to the suppressors of suppres
                  trehalose-6-phosphate phosphatase,
                  as well as propos., in particula:,
                                                                                                                                        * Marical prepas., which include
                  innibitors or suppressors obtained to
                                                                                                                                        screening
                 assay. Inhibitors are described .
                                                                                                                                               es applications in
                 biocides and antifungal pharmage of
               9023-07-8, Sugar phosphatase 9025-72-3
                   , Trehalose-6-phosphate phosphatase
                 9055-29-2, Mannitol-1-phospharas 37228-75-4,
                  Glycerol-3-phosphatase
                 RLE BAC (Biological activity or Htt. ) , Hwoept adverse; BER (Biological
                 process'; BSW Biological study, .:
                                                                                                                                           cred ; BILL Biological study;
                 PROC Process'
                          inhibitors; sugar alc. phosp::// sugar
phosphatases as novel targets : antiparasitic
                          agents and use of inhibitors . broordes and pharma way reals
 L19 ANSWER & OF 15 HOAPLUS COPYRIGH ACCESSION NUMBER: 2000:742235 at 2000MENT NUMBER: 133:291952
                                                               Modification of colosynthesis by DNA shuffling Yuan, Ling; Fill of Sun Ai; Lassner, Michael Maxygen, Inc., FCT Int. App.,
   TITLE:
   INVENTOR(S):
  PATENT ASSIGNEE(S):
 SOURCE:
                                                                            CODEN: PIXXD.
 DOCUMENT TYPE:
                                                                            Patent
  LANGUAGE:
                                                                             English
  FAMILY ACC. NUM. COUNT: 1
  PATENT INFORMATION:
              PATENT NO. KIND DATE WILLICATION NO. DATE

WO 2000061740 A1 20001019 WILLICATION NO. DATE

W: AE, AL, AM, AT, AU, AZ, FA, S, S, BE, BY, CA, CH, CN, CR, CU, CZ, DE, DW, DM, EE, ES, FL, S, L, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, FF, S, C, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MI, S, L, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TL, S, S, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TI,
                            SK, SL, TJ, TM, TR, TT, T., AZ, BY, KG, KZ, MD, RU, T., RW: GH, GM, KE, LS, MW, SD, FI,
RW: GH, GM, HE, LS, MW, SD, F1, ..., II, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, II, ..., MC, ML, PT, SE, BF, BU, CF, CG, CI, CM, GA, GN, SW, MI, II, E, SN, TD, TG
PRIORITY APPLN. INFO.:

AB Methods of modulating lipid prodm. Shadow whole organisms by DNA shuffling are provided. Single gets a surrous limit biography.
               shuffling are provided. Single q_{\rm eff}, regrons, lipid biosynthetic cycles and whole genomes can be recombined as graduate cells and organisms with
                desirable lipid synthetic or metal and invity. Libraries of recompined
                lipid synthetic nucleic acids and the ms are also provided.
              Modification of lipid sath, fathy compn., acyl chain length, located specific to the described. A decrease in susception of methyltransforase in compn., acyl chain, cyclopropyl, method chain, cyclopropyl, method specific described. Use of two-hybrid system to many the chain activity is also also also screening to libraries,
```

```
such as phage display library is a such as sorn, peanut, parley, millet, rice, soyned and such as sorn, nut whise lipid prosynthetic anticle and short and short as a powerful process of the source o
                 Jenes.
  REFERENCE COUNT: 11 THERR AND FOR REFERENCES AVAILABLE FOR THIS BECOME. OF US AVAILABLE FOR THIS
  119 ANSWER TO IF IS HOAFLUS COPYRIDG.
  ACCESSION NUMBER: 100018088000
                                                                     The non-meval: promited pickynthesis of plants
                                                                    as a test symbol who now herbicides and
                                                                    drugs against : . . : hacteria and the
                                                                   malaria parasite
  AUTHOR S :
                                                                    Lightenthaler, waster M.; Deidler, Johannes;
                                                                Schwender, Cristian, Christian, Botanisches Immarker II, Universität Karlsruhe, Karlsruhe, Immany
  CORPORATE SCUPPE:
 P. falciparum, and cures malaria-th: " : mide. This is the first
              successful application of a herbinia of the novel isoprenoid pathway as a
              possible drug against malaria.
 REFERENCE COUNT: 40 THERE ARE 1 FOR PREFERENCES AVAILABLE FOR THIS RECURLOUS AVAILABLE IN THE RE FORMAT
  L19 ANSWER 8 OF 15 HCAPLUS COPYRIGHT
 ACCESSION NUMBER: 2000:6150 H WA DOCUMENT NUMBER: 132:307079
TITLE: Characterisar. 10.0 % % fit gene licated between the class II report 10.0 % genes in the numan MED AUTHOR'S: Aguado, E., 10.0 % F. I.

CORRORATE SOURCE: MEC Immunities 10.1, Switzra University, Owford, OXI 300, UF
                                                                   SOURCE:
```

```
325 1. ", "I+ Fale +
                                                                                                           Histodompatin. Son pland Tinierence], 12th, Saint-Main and Senance, 1294 [1947], Meeting Late 1266, Vily Senance, Lett. Land Committee in Thermatical Committee in the Committee
                                                                                                            Publisher: 1000
                                                                                                             OLĪEN: FEMBA
   DI MUNEUT TYPE:
                                                                                                             Conference
     LANGUAGE:
                                                                                                          Enalish
  AB The navel gene GIS encodes a les as a sprometr with a predicted mol. with of about 32 kDa which contains and a transference segments. The GIS gene is a single copy gene, the contains of t
                       cells. The protein shows hemol. w. - *
                                                                                                                                                                                                enzyme LFAAT
                           l-acyl-sm-glycerol-3-phosphate
                     bacteria. The authors expressed 3 insect cells insect calls assays whether Glb is the number 3: to deminstrate by encymber localization of the enzyme.
   REFERENCE COUNT:
                                                                                                        10 THERE ARE TO REPRENCES AVAILABLE FOR THIS RECORD. FOR SECURION AVAILABLE IN THE RESENTANT
 L19 ANSWER 9 OF 15 HCAPLUS CURYRIGHT ACCESSION NUMBER: 1999:775162 H DOCUMENT NUMBER: 132:148891
                                                                                                        Determination trehalose-6-
phosphate leve antharomyces derevisiae,
using Bacillus and as phosphotrehalase
   TITLE:
                                                                                                      Van Vaeck, C.; L.:., I.; Bonini, B.; Van Dijck, P.;
Thevelein, J. M.
Laboratory of M. Gover Cell Biology, Institute of
Botany and M. St. Governor, Louvain, Belg.
Mededelingen - Governor, Conversiteit Gent; (1999),
 AUTHOR (S):
  CORPORATE SOURCE:
 SCURCE:
                                                                                                        64(5b), €47---
                                                                                                        CODEN: MFLBER; ".: 1473-7503
                                                                                                        Universiteit Ders, Frühreit Landbouwkundige en Toegepaste B.W. Wetenschappen
 PUBLISHER:
  LOCUMENT TYPE:
                                                                                                          Journal
  LANGUAGE:
                                                                                                         English
 AB The disaccharide trehalose is a manage of the stress resistance in
                     several organisms. It was found in hacteria, yeast,
                     fungi and in certain invertebrath \pi . It species. Deletion of
                     the 1st enzyme of trehalose synthem, . . . . in yeasts results in a pleiotropic phenotype inclusion as the of trehalose, deficiency in
                     growth on rapidly fermentable sugar. ...: .:ss of stress resistance. To
                     study the changes of trehalose-6-phosphate
                      (Tre6P), the precursor of trehalist, + 0.00 type cells and in yeast cells
transformed with TPS1 homologs from the regardisms, the authors developed a novel Tre6P assay. The authors' to the show that complementation of a tps1.DELTA. Str. in homologs from other organisms restores growth, but not proper that the sugar influx into glycolysis.

REFERENCE COUNT:

5 THERE ARE TO REFERENCES AVAILABLE FOR THIS RECORD. A CATI MS AVAILABLE IN THE RE FORMAT
 LIP AMSWER IS OF IS HOAPLMS COPYRIGHT
ACCESSION NUMBER: 1999:464099 E
 DOCUMENT NUMBER:
                                                                                                         131:94934
                                                                                                  Flan' galar ...
                                                                                                                                                                                                     14 N.1848
```

i, Heir

Smirniii, Wilk;

INVENTOR'S :

```
946 17 207, 314 Page 9
 FATENT ASSIGNED S: ASCITAGE Limited (1) to technology Courawest Limited SIGNOR: POT Int. App...
  1.0WMENT TYPE:
                                                                                Faterit
 IANGUAGE:
FAMILY ACC. NUM. COUNT:
FATENC INFORMATION:
                                                                                Er. 911281.
              PRIORITY APPLN. INFO.:
           This invention presents and charge an enzyme which catalyzes the converse of L-galactose dehydrogenase, an enzyme which catalyzes the converse of L-galactose to L-galactose is an oxidi. The catalytic conversion conversion is an oxidi. The catalytic conversion conversion is an oxiditation of the catalytic conversion catalytic conversion catalytic conversion catalytic conversion conver
                a herbicidal compn. comprising a new contract inhibits L-galactose
                dehydrogenase.
                                                                                                THERE ARE COMES REFERENCES AVAILABLE FOR THIS RECOPD. A CONTIONS AVAILABLE IN THE RE FORMAT
REFERENCE COUNT:
 119 ANSWER 11 OF 15 HOAPLUS COPYRIGHT
ACCESSION NUMBER: 1998:246371 HT
DOCUMENT NUMBER:
                                                                                129:64817
                                                                            Trehalose-6-phosphate
                                                                            phosphatases is a complementation of the yeast tps2 minutes.
                                                                         AUTHOR(S):
COPPORATE SOURCE:
                                                                              CH-4116, Switz.
```

```
345 - 1 171, 11<del>5</del> - Page 11
                                       SIMBOE:
 1 11 1 2 11 12 1
 L. DIMENT TYPE:
                                        Jagraal
                                        English
AB It is currently thought that most in the plants lack the dapacity to synthesize trehalose, a common dost to the city a major role in desideation tolerance. Attempts to the fire liven made to render plants more drought-resistant by the excited process to increase the synthesis. It is demonstrated not processed genes for at least one in engages required for trehalose synthesis.
         trehalose synthesis, trehalose-6-phosphate
        phosphatase. The yeast tysl mutahr, and lacks this enzyme, is heat-sensitive, and Arar and a complement
        enzyme, is heat-sensitive, and Aramara, commander this effect has been screened for a command the peast
        transformants that grew at 36.6.00: It also able to produce trenalose. All of these empresses the Arabidopsis office, either AtTREA or AtTREE, which are bosts to the C-terminal part of the
        yeast TFS2 gene and other migra; ... trehalose-6-
        phosphate phosphatases. Weast the second expressing
        AtTPPA or AtTPPB contained trehalose 6
        both in vivo and in vitro. The enzyme and sphorylated
        trehalose-6-phosphate but not
        glucose-6-phosphate or sucrose-\epsilon-\epsilonn \rightarrow \epsilon. Both genes are expressed in flowers and young developing tissue \epsilon and appears. The finding of these
        novel Arabidopsis genes for trehalose-6 -
        for trehalose biosynthesis exists at the same.
        9025-72-3, Trehalose-6-phosphate
        phosphatase
        RL: PRP (Properties)
              trehalose-6-phosphate
             LIP AMSWER 12 OF 15 HOAPLUS COFYROUSE
ACCESSION NUMBER: 1993:230685 HT
DOCUMENT NUMBER:
                                        118:239685
                                    Effect of engines of a bacteria on mitochondrial - provide activities in the weevil Sitophilus ory: A properties: Curculionidae) Heddi, A.; Lefect of the lyon, F. Lab. Biol. Applies to the lyon, Villeurbanne, 69621, Fr. Insect Biochem.
TITLE:
AUTHOR(S):
CORPORATE SOURCE:
SCURCE:
                                      23(3), 403-11
                                       CODEN: IEMBES; CODEN: 465-1748
DOCUMENT TYPE:
                                       Journal
LANGUAGE:
                                       English
AB Various mitochondrial enzymic actives a were investigated in symbiotic and aposymbiotic larvae and adults of gallus cryzae. Six enzymes were assayed: cytochrome actives, succinate
        sytochrome o reductase, glycerol 3-phosphate
       cytochrome o reductase, iscoltrate to a mase, pyruvate dehydrogenase, and .alpha.-Retoglutarate denydrome to the specific activities of all these enzymes were higher in mit to the specific activities of all symbolic larvae than those iscoltrate to a graymrictic larvae. In adults, the differences in encymic activities to sympositic and aposymplicities
```

```
- 1 ., 1- Fage 11
                         Flant Sturnal 1994 , 13 : , 677-645 
Silen: Floven; ISSN: 1961-7411
Blankwell Aniembe Dha.
Journal
                         English
. Ai. A E:
    in it carrently to aght that mist fliwering plants lack the capabily o
     Contresize trenal.se, a common disacon mide : bacteria,
     fungi and unvertexistes that appears to play a magor role in
     designation tolerance. Attempts have therefore peen rade to removing lands
     non-grought-resistant by the expression of mirrobial genes for trenalise
     synthesis. It is demonstrated here that Arabidopsis thalland itself
     possesses jenes for at least one of the enzymes required for
     premainse synthesis, trehalose-6-phosphate
    phosphatase. The yeast tysl mutant, which lakes this
     enzyme, is heat-sensitive, and Arabidopsis olNA above to complement
     this effect has been screened for. Half of the yeast
     transformants that brew at iffice byron. A work enact which type disk
     trenal so. All i these expressed the i two Aracinists (INA, ectner
     Attifia or Attifie, which are both nomologous to the 2-tyrminal part of the
     yeart TFSz gene and other microbial trehalose-6-
     phosphate phosphatases. Yeast tps2 mutants expressing
     AtTEFA or AtTPPE contained trehalose-6-
    phosphate phosphatase activity that could be measured
     noth in vive and in vitro. The enzyme dephosphorylated
     trehalose-6-phosphate but not
     glubosé-6-phosphate or suprose-6-phosphate. Both genes are empressed in
     flowers and young developing tissue of Arabidopsis. The finding of these
     navel Arabidopsis genes for trehalose-6-
     phosphate phosphatase strongly indicates that a pathway
     for trehalose biosynthesis exists in plants.
    9025-72-3, Trehalose-6-phosphate
    phosphatase
     RL: PRP [Properties,
         trehalose-6-phosphate
        phosphatases inom Arabidopsis thaliana: identification (y
        functional complementation of yeast tps2 mutant
LIP AMSWER 12 OF 15 HOAPLUS COPYRIGHT 2003 ACS
ACCESSION NUMBER:
                         1993:230685 HCAPLUS
                          119:230685
DOMINENT NUMBER:
Effect of endocytobiotic bacteria on
                         mitochondrial enzymic activities in the weevil
                         Sitophilus oryzae (Coleoptera: Curculion: dae,
                         Heddi, A.; Lefebvre, F.; Nardon, F.
Lab. Biol. Appl., INSA Lyon, Villeurbanne, 69621, Fr.
Insect Biochemistry and Molecular Biology (1998),
ATTHER S':
CLEFIFATE SOURCE:
BYTH E:
                         23(3), 403-11
CODEN: IBMBES; ISSN: 0965-1746
                          Journal
I THE
LAN MAGE:
                         English
     Various mitophondrial engymic activities were investibated in symblotic
     and appropriate darvae and adults of Sitophilus oryvae. Tim
     enzymes were assayed: syttochrome a swidase, succinate
     nyto mrome o reductase, glycerol 3-phosphate
     synthing on reductase, isositrate dehydrogenase, pyruvate whyso senase,
     and lappha. - Retoglistarate denymit genase. The specific activities of all
     these enzymes were nigher in mittahandria isolates in m
     symmittic larvae than thise isilated from apisymmit of Laften. The state,
     the differences in encymia activities between symbolic and district to
```

```
insects were altendated. .alpha. - Februar ate buyur becase
           aminity was similar in the Distrains, and pyrimate benying sender aminity, who have the appropriate timestrain. From the results of the sense and
           Carming sees telement sympletic and aposymilation insects in
           Althors of the that the present of bacteria is responsible for the plants and the present of bacteria is responsible surface for the plants enough that the present of bacteria. These demoys a strength of the mit of the properties of the propertie
           is later from Larval bacteriome. We approvity was upon
  II - ANUMER IS OF IE HOAFLUS COPYRIGHT LICE ACS
  ADDROSION NUMBER:
                                             1972:174398 HOAFLTS
                                                    116:104394
                                                  Target gene-complemented mior organisms for identification of antiparasite drugs Miein, Bonald D.; Greary, Fin thy S.
  1117811718 8 :
 PARENT ASSIGNEES:
                                                  Tprohn Co., USA
POT Int. Appl., Dé 55.
CODEN: FIXMES
 Name E:
 Fateri
 Emalish
 PARTLY AND. NUM. OUTST:
 PACKET INFORMATION:
           PATENT NO. KIND DATE
                                                                                      APPLICATION NO. LATE
           -----
                                                                                      ----------
           WU 9007260 A1 19900014
                                                                                    WU 1991-V82707 (1991.428
                  WE AU, BB, BB, BB, CA, FI, BU, JP, KP, KB, LK, MM, MB, MW, M , BI,
                  FO, SI, ST, TS

EW: AT, BE, BF, BJ, CF, CG, CH, CM, DE, DK, ES, FR, GA, GB, GR, IT,
                        LU, ML, MR, NL, SE, SN, TD, TG
          US 8179143
AU 9179750
                                                                                      US 1990-517633
                                                                                                                        19900502
                                                        19911127
                                                                                     AU 1991-79750
                                                                                                                         19910425
                                                                                US 1990-510633
WO 1991-US2067
 PRICEITY APPLN. INFO.:
                                                                                                                         19900502
                                                                                                                         19911428
       A method for identifying antiparasitic drugs comprises exposing
          parasite gene-complemented microorganisms to the test
          compd. and detg. microbial viability. An Escherichia coli mutant
          deficient in both phosphofructokinase (FFK) enzymes was used to
          plane the PFW pDMA of Haemonphus contortus by complementation. The SDMA
          was sequenced.
119 AMSWER 14 OF 18 HOAPLUS COPYRIGHT 2003 ACS
ACCESSION NUMBER:
                                           1988:90393 HCAPLUS
1000MENT NUMBER:
                                                   108:90898
                                                 A sensitive and efficient iscentione teamnique for
                                                 small arthropods and other invertebrates
ATTERIE
                                                 Easteal, Simon; Boussy, Jan A.
                                                  Pas. Soh. Biol. Coi., Aust. Mari. Thir., Canteerra,
 THE RATE SOURCE:
                                                  2611, Australia
JOSEPH E.
                                                  Bulletin of Entomological Research 1987,, " 3 ,
                                                  407-15
                                                  COUEN: BEREAL; ISSN: 0000-4883
                                                  Journal
LAN PAGE:
                                                 Er ;lish
Ab . An electrophyretic method for the study of enzyme varietion,
         which uses cellulise abotate shoots with an agar operlay for staining, the
         use of a very good general purpose buffer "bitrib-aminopropylaistnan".
          write, and the use of sodium addite as a bacteriocide the allow-
```

```
. Di term storade (i dhems. às solhs. are resonired. Tests are
           reported in the technoque on Tetrany mus introde Form, Aedes aegypti, and
           seberal species in Trisophila. The temnique offers sensiolving equal t
            is dreater than staith or polyadrylarine get electrophiceeus and i-
           specialist to very small manistrs, allowing either the testing
            is not the individuals in the hos. It enzymes is the
           testing I lewer enzymes distribution of translations and the continues of 
           the termingue is emitypent with respect to time and materials, and safer
           Than I have the half methods.
          9075-65-4, Glycerol 3-phosphate
           Hong in private
          Blacklif Billian at study
                  istendymes, detection of, by deligious adetate electronic rests
 Dis Anomer de un 18 HOAPLUS CORMEDSHT LUIS ADA
A MESSI D DUMBER: 1968:1968: HOAPLU
A MESMI N NUMBER:
                                                  1:114868
                                                 Effects of some antitumor agents on growth and
                                                 glycolytic enzymes of the flagellate
                                                  <u>Örithidia</u>
                                                Bacchi, Cyrus J.; Ciaccic, Edward I.; Koren, Luis E. Haskins Lab., New York, NY, USA
Journal of Bacteriology (1969), 96 1 , 23-6
COZEN: JOBAAY; ISSN: (221-8193
AUTHUR \varepsilon:
 CURPURATE SOURCE:
S TROBE
Journal
LANGERE:
                                                 English
Ah - Some antitumor agents Anown to specifically inhinit pertain tumor sel.
         enzymes were examd. for activity against glycolymi:
         enzymes and for growth of the insect trypanosomatid, C.
          fasciculata. The cytoplasmic enzymes hexckinase,
         .alpha.-glycerophosphate dehydrogenase, malic dehydrogenase, and
          glusose-6-phosphate dehydrogenase were tested. Agaricio acid
          .z-hydroxy-1,2,3-nonadecanetricarboxylic acid: was nighly inhibitory
         50-100 ) to malic and .alpha.-glycerophosphate dehydrogenases at .aptrm.s..times. 10-6M; 2-.p-hydroxyphenyl:-2-phenylpropane (0..times. 10-4M) and 6,6-aichlurc-2-penzomazolinone (5..times. 10-4M) were less effective >50.
         inhibition, against them. The antiprotozoal agents primaquine ,4 .times.
         10-4M) and Melarsoprol (8 .times. 10-4M) were 30-40% inhibitory. Agarists
         acid, 2-(p-hydroxyphenyl)-2-phenylpropane, and 5,6-dichloro-2-
         bencomagolinone inhibited growth of Crithidia at less than 10-4M. Elight
         other test compds. from the Cander Chemotherapy National Service
          Center (CCNSC) were not toxic to dell growth, although two
          (4-biphenyloarbowylic abid and 1-(p-chloropenzyl)-2-éthyl-5-methylindole-3-
         abetin acid) inhibited Crithidia .alpha.-glyberophosphate dehydrogenase
below 1M. A.1 of the compds. used specifically inhibited canber bell
         .alpna.-glycerophosphate dehydrogenase. The corresponding enzyme
         in pathogenic African orypanosomes is important in their termina.
         respiration. C. fasciculata may pequeefil in preliminary evaluation to
         chemitherapeutic agents as potential trypanoribus.
        9075-65-4, G.y erol prosphate denydrogenase
                in Crithidia fasciculata, neoplasm inhibitor eirest on
```